



50




REVIEWS in
MINERALOGY &
GEOCHEMISTRY

Volume 50



— *BERYLLIUM* —
MINERALOGY, PETROLOGY,
AND GEOCHEMISTRY

EDWARD S. GREW, EDITOR



MINERALOGICAL SOCIETY OF AMERICA

Paul H. Ribbe, *Series Editor*

GEOCHEMICAL SOCIETY

Jodi J. Rosso, *Series Editor*

— *BERYLLIUM* —
MINERALOGY, PETROLOGY, AND GEOCHEMISTRY

G
MSA

CONTENTS

- 1 **Mineralogy, Petrology and Geochemistry of Beryllium: An Introduction and List of Beryllium Minerals** *Edward S. Grew*
- 77 **Behavior of Beryllium During Solar System and Planetary Evolution: Evidence from Planetary Materials**
Charles K. Shearer
- 121 **Trace-Element Systematics of Beryllium in Terrestrial Materials**
Jeffrey G. Ryan
- 147 **Rates and Timing of Earth Surface Processes From In Situ-Produced Cosmogenic Be-10**
Paul R. Bierman, Marc W. Caffee, P. Thompson Davis, Kim Marsella, Milan Pavich, Patrick Colgan, David Mickelson
- 207 **Cosmogenic Be-10 and the Solid Earth: Studies in Geomagnetism, Subduction Zone Processes, and Active Tectonics**
Julie D. Morris, John Gosse, Stefanie Brachfeld, Fouad Tera
- 271 **Environmental Chemistry of Beryllium-7**
James M. Kaste, Stephen A. Norton, Charles T. Hess
- 291 **Environmental Chemistry of Beryllium** *J. Veselý, S.A. Norton, P. Skřivan, V. Majer, P. Krám, T. Navrátil, J.M. Kaste*
- 319 **Beryllium Analyses by Secondary Ion Mass Spectrometry**
Richard L. Hervig
- 333 **The Crystal Chemistry of Beryllium**
Frank C. Hawthorne, Danielle M.C. Huminicki
- 405 **Mineralogy of Beryllium in Granitic Pegmatites** *Petr Černý*
- 445 **Beryllium in Silicic Magmas and the Origin of Beryl-Bearing Pegmatites** *David London, Joseph M. Evensen*
- 487 **Beryllium in Metamorphic Environments (Emphasis on Aluminous Compositions)** *Edward S. Grew*
- 551 **Be-Minerals: Synthesis, Stability, and Occurrence in Metamorphic Rocks** *Gerhard Franz, Giulio Morteani*
- 591 **Non-pegmatitic Deposits of Beryllium: Mineralogy, Geology, Phase Equilibria and Origin** *Mark D. Barton, Steven Young*

ISBN 0-939950-62-6



9780939950621

Environmental Chemistry of Beryllium-7

James M. Kaste

*Department of Earth Sciences
Dartmouth College
Hanover, New Hampshire 03755
james.m.kaste@dartmouth.edu*

Stephen A. Norton

*Department of Geological Sciences
University of Maine
Orono, Maine 04469*

Charles T. Hess

*Department of Physics and Astronomy
University of Maine
Orono, Maine 044699*

INTRODUCTION

In addition to the stable isotope ^9Be , Be is also formed as two cosmogenic isotopes of interest to earth scientists. Cosmogenic Be is formed primarily in the stratosphere from cosmic-ray spallation of oxygen and nitrogen, but some is produced in the troposphere and *in situ* on the surface of the earth. Production of cosmogenic Be was hypothesized by Peters (1955). Soon after, naturally occurring ^7Be was identified in precipitation by Arnold and Al-Salih (1955) and essentially concurrently and independently by Goel et al. (1956). ^{10}Be ($T_{1/2} = 1.5 \times 10^6$ yr) was first detected by Arnold (1956) in marine sediment cores. The $^{10}\text{Be} / ^7\text{Be}$ production ratio is approximately 0.5 (Lal and Peters 1967).

After its formation in the atmosphere, cosmogenic Be adsorbs electrostatically to aerosols that may be washed out by precipitation and delivered to ecosystems. The amount of cosmogenic Be that reaches the surface of the earth is a function of production rate (cosmic-ray intensity), stratosphere-troposphere mixing, circulation and advection within the troposphere, and efficiency of removal from the troposphere (wet and dry deposition) (Feely et al. 1989). Since the discovery of cosmic-ray produced Be, numerous researchers have gathered data on its (1) production, (2) fluxes to the oceans and terrestrial ecosystems, (3) distribution and inventory in soils, snow, sediments, and vegetation, and (4) geochemical behavior. In 1955, Peters suggested that the relatively long half-life of ^{10}Be could make it useful for quantifying Tertiary sedimentation rates and other surficial processes. The focus of this chapter is the short-lived radionuclide ^7Be ($T_{1/2} = 53.12 \pm 0.07$ days; Jaeger et al. 1996). Due to its short half-life, relative ease of measurement, and well-defined source term, ^7Be serves as a useful tool for tracing and quantifying environmental processes on the <1 year timescale (Lal et al. 1958; Young and Silker 1980; Krishnaswami et al. 1980; Russell et al. 1981; Turekian et al. 1983; Dutkiewicz and Husain 1985; Olsen et al. 1985; Wallbrink and Murray 1993; Bonniwell et al. 1999), and has applications in meteorology, soil science, sedimentology, geomorphology, hydrology, geochemistry, and nuclear physics.

ANALYSIS FOR ^7Be

^7Be decays to stable ^7Li by electron capture ($\lambda = 4.766 \text{ yr}^{-1}$). Eighty-nine and one-half percent of the atoms decay directly to the ground state of ^7Li , while 10.5% decay first to the excited state of ^7Li , which decays to ground state ^7Li via gamma-ray emission at 477.6 keV (Ajzenberg-Selove 1988). ^7Be activities are normally determined in

environmental samples using gamma spectrometers that detect the 477.6 keV gamma. In earlier investigations, ^7Be was detected using thallium (Tl) activated sodium-iodide (NaI) detectors (Arnold and Al-Salih 1955; Goel et al. 1956; Walton and Fried 1962; Schumann and Stoeppler 1963, and Krishnaswami et al. 1980). However, the relatively low resolution of these instruments requires chemical separation of the Be before the gamma analysis because the NaI(Tl) detector cannot distinguish the ^7Be photopeak from other radionuclides which decay in the same energy region (^{228}Ac at 462 keV; ^{103}Ru at 497 keV). The techniques used for the purification and concentration of ^7Be from various samples generally utilize cation exchange columns and/or coprecipitation of Be with Fe hydroxides and other hydroxide phases. Details of these methods are described in the references mentioned above.

More recently, high-purity germanium (Ge) detectors or lithium (Li)-drifted Ge detectors have been employed to determine ^7Be activity. These detectors generally have high resolution (<2 keV) and high counting efficiencies. In many cases, precipitation, soil, and sediment samples can be analyzed directly for ^7Be . Larson and Cutshall (1981) describe a method for detecting ^7Be directly in sediment samples using Marinelli Beakers. They reported a detection limit of approximately 0.015 Bq g^{-1} for a 250 g sediment sample counted 200 min ($1 \text{ Bq} = 1 \text{ becquerel} = 1 \text{ disintegration per second}$, the SI unit for radioactivity. $1 \text{ Bq} = 27.03 \text{ pCi}$; no concentration equivalent such as ng kg^{-1} or pg kg^{-1} would be applicable due to the very small number of atoms present in environmental samples). When counting samples high in ^{232}Th , one should be cautious of the 478.3 keV decay energy of ^{228}Ac . Although the relative intensity of this photon is only 0.215%, high activities of ^{228}Ac found in mineral soil and sediment samples can create a disturbance in this region of the spectrum. Surface waters are generally low in ^7Be , so pre-concentration is necessary for efficient detection. Kostadinov et al. (1988) suggested ^7Be adsorption on activated charcoal to concentrate ^7Be from surface waters.

For over five decades, ^7Be has received attention from scientists investigating the possible variability in a radioactive nuclide's decay constant. Emilio Segrè (1947) first suggested that the decay constant of a nuclide that decays by electron capture should be influenced by the density of its electrons. He suggested that, for a light element such as ^7Be , this change could be "appreciable", and by putting ^7Be in various chemical compounds, a slight change in half-life could potentially be observed. A few years later, Leininger et al. (1949; 1951) reported that $[\Delta\lambda/\lambda] \times 10^{-3}$ for $\lambda(^7\text{Be}) - \lambda(^7\text{BeF}_2)_{\text{hex}} = 0.84 \pm 0.10$. Between 1950 and 1970, several researchers reported a deviation of λ for ^7Be in different Be compounds (Table 1). These reports received little attention from the scientific community because the differences in λ were detected using NaI detectors or ionization chambers. The analytical error of these instruments approached the magnitude of changes in λ they were trying to resolve. In 1999, using a high purity intrinsic Ge detector, Huh (1999) reported that the half-lives of ^7Be in $\text{Be}(\text{OH}_2)^{+2}$, $\text{Be}(\text{OH})_2$, and BeO are 53.69, 53.42, and 54.23 d ($\pm 0.1\%$), respectively. Changes in the chemical form of ^7Be may not be the only variable affecting its decay constant. Hensley et al. (1973) found that the decay constant of ^7Be in BeO increased slightly under high pressure in a diamond-anvil press. Liu and Huh (2000) found similar results as Hensley et al. (1973), but they noted that the rate of increase of λ with pressure in a $\text{Be}(\text{OH})_2$ gel decreased with increasing pressure.

PRODUCTION AND DELIVERY OF ^7Be TO THE EARTH'S SURFACE

Production of ^7Be depends on the cosmic-ray flux, which varies with latitude, altitude, and solar activity. Lal et al. (1958) calculated production rates for cosmogenic Be. Production in the troposphere increases by a factor of approximately 1 to 3 from the equator to the poles. In the stratosphere, production can increase by a factor of 4 to 5

from the equator to the poles. Production is greatest between 12 and 20 km altitude (depending on latitude), decreasing nearly exponentially by about 3 orders of magnitude to the earth's surface (Lal et al. 1958; Bhandari et al. 1970). Cosmogenic Be production varies with the 11-year solar cycle. Solar activity maximums result in increased deflection of cosmic rays from the solar system (Lal and Peters, 1967) that decreases the cosmic-ray flux to the earth, and thus decreases ^7Be production. Several authors have demonstrated an inverse relationship between cosmogenic Be concentrations in air and on the surface of the earth with solar activity (Fig. 1) (Beer et al. 1990; Hötzel et al. 1991; Durana et al. 1996). Concentration fluctuations in surface air resulting from the 11-year solar cycles are generally on the order of 15 to 25% (Koch and Mann 1996) (Fig. 1).

Table 1. Variation in the decay constant of ^7Be in various chemical compounds. All values are $\Delta\lambda/\lambda \times 10^{-3}$. Modified from Jöhlige et al. (1970).

Source pair	Leininger et al. (1949); Segrè & Wiegand (1951)	Kraushaar et al. (1953)	Bouchez et al. (1956)	Jöhlige et al. (1970)	Huh (1999)
$\lambda(\text{BeO}) - \lambda(\text{BeF}_2)_{\text{hex}}$	0.69±0.03	0.609±0.055			
$\lambda(\text{BeO}) - \lambda(\text{BeF}_2)_{\text{am}}$				1.130±0.058	
$\lambda(\text{Be}) - \lambda(\text{BeF}_2)_{\text{hex}}$	0.84±0.10	0.741±0.047			
$\lambda(\text{Be}) - \lambda(\text{BeF}_2)_{\text{am}}$			1.2±0.1		
$\lambda(\text{BeO}) - \lambda(\text{Be})$	-0.15±0.09	-0.131±0.051			
$\lambda(\text{BeO}) - \lambda(\text{Be}(\text{OH})_2)$					-15.177±0.1




Figure 1. Annual variation of ^7Be in surface air plotted with solar activity, 1955-1993. ^7Be units are millibecquerels m^{-3} at standard temperature and pressure. Figure used by permission of the editor of *Journal of Geophysical Research*, from Koch et al. (1996), Fig. 2, p. 18,653.

Because the stratospheric residence time of aerosols (approximately 14 months [Reiter, 1975]) exceeds the half-life of ^7Be by more than six-fold, equilibrium activity is assumed to occur. In the troposphere, however, production of ^7Be is significantly lower, and the residence time of ^7Be is much shorter (22 to 48 days [Bleichrodt 1978; Durana et al. 1996]) due to rapid wash-out. This results in a fairly high concentration gradient between the stratosphere and the troposphere, with tropospheric air generally containing one to two orders of magnitude less ^7Be (Bq m^{-3}) than the stratosphere. Rama and Honda (1961) and Bhandari and Rama (1963) reported ^7Be activity for air sampled at various points in the atmosphere. Stratosphere air sampled at 18-20 km ranged from approximately 0.16 to 0.58 $\text{Bq } ^7\text{Be m}^{-3}$. Troposphere air ranged from approximately 0.005 to 0.02 Bq m^{-3} . Dutkiewicz and Husain (1985) found that ^7Be activity in lower stratospheric air averaged approximately 0.17 Bq m^{-3} , while the upper troposphere averaged 0.02 Bq m^{-3} .

While the activity of ^7Be in the stratosphere remains fairly constant, concentrations in the troposphere and near-surface air, and thus the amount of ^7Be available to ecosystems exhibit seasonal fluctuations (Feely et al. 1989). ^7Be concentrations in near-surface air generally range from 0.001 to 0.007 Bq m⁻³ (Feely et al. 1989; Dueñas et al. 1999). Stratosphere-troposphere exchange can increase ^7Be concentrations in the troposphere and near-surface air. Husain et al. (1977) correlated high concentrations of ^7Be at Whiteface Mountain, New York, USA with stratospheric air masses as indicated by potential vorticities. Viezee and Singh (1980) correlated high concentrations of ^7Be in surface air with low-pressure troughs. Maximum mixing between the stratosphere and the troposphere generally occurs in the spring of each year at mid-latitudes, and near-surface air at the middle latitudes generally has higher concentrations of ^7Be at this time (Feely et al. 1989). Intense thunderstorms may also mix stratospheric air downward, thus increasing the amount of ^7Be available for scavenging by precipitation (Dingle 1965; Noyce et al. 1971). Several authors have suggested that vertical mixing within the troposphere also governs the concentration of ^7Be in rain and near-earth surface air (Feely et al. 1989; Baskaran 1995; Durana et al. 1996; Dueñas et al. 1999). Warming of the earth's surface during the spring and summer will increase convection, which would transport ^7Be from the upper troposphere to near-surface air. At latitudes >60°, seasonal variations in surface air result from the transport poleward of mid-latitude air. Arctic sites commonly have peak ^7Be air concentrations in the late winter or early spring, probably a result of the arrival of mid-latitude tropospheric air high in ^7Be (Feely et al. 1989). Sites with a pronounced seasonal variation in rainfall amount show an inverse relationship between rainfall rate and ^7Be concentration in air, which demonstrates that washout of ^7Be can have a significant impact on surface air concentration (Feely et al. 1989). Koch and Mann (1996) used a singular value decomposition (SCD) technique to analyze the spatial and temporal variability of near-surface air concentrations of ^7Be . They found that up to 47% of the monthly air concentration variability could be explained by 4 "modes" which correspond to the 11-year solar cycles, El Niño Southern Oscillation cycles (2 to 7 years), annual, and semi-annual cycles. Zanis et al. (1999) measured ^7Be at an alpine research station in Switzerland from 4 April 1996 to 1 January 1997, and compared this with meteorological parameters and composite 500 hPa geopotential height maps. They concluded that downward transport of air from the upper troposphere and wet scavenging of aerosols were main processes controlling the ^7Be concentrations measured at the site. Papastefanou (1991) found that a supernova explosion in 1987 increased the ^7Be activity in air by up to four times the normal activity found in temperate air masses. Nuclear detonations do not contribute to ^7Be production.

Arnold and Al-Salih (1955) were the first to report cosmic-ray-produced ^7Be in wet precipitation. They collected large volumes of rain and snow in Indiana and Illinois from October 1953 until April 1954. ^7Be was concentrated from the samples by adding a stable Be carrier and co-precipitating Be(OH)₂ along with Fe(OH)₃ and other hydroxides at a pH of 9. The precipitate was then purified to contain only BeO and counted using a 2.54 cm NaI (Tl) crystal scintillation spectrometer with an overall counting efficiency of 0.99%. ^7Be activities ranged from 0.16±17% to 2.40±8% Bq L⁻¹, with a volume-weighted mean of approximately 0.95 Bq L⁻¹. While the Arnold and Al-Salih (1955) paper was in press, Goel et al. (1956) independently demonstrated the presence of ^7Be in Bombay, India. During the months of July, August, and September of 1955, they documented ^7Be in monsoon rain with a volume-weighted mean activity of 0.36 Bq L⁻¹. Since then, numerous researchers have reported the activity of ^7Be in wet-precipitation, with annual volume-weighted means generally ranging from 0.75 to 2.75 Bq L⁻¹. Baskaran et al. (1993) tabulated the volume-weighted ^7Be activities in precipitation reported between 1983 and 1993. Some studies prior to 1980 did not acidify their precipitation samples during collec-




Figure 2. Seasonal variation of ⁷Be flux at two sites. Data plotted from Olsen et al. (1985).

tion, which may have resulted in poor recovery of the ⁷Be from the precipitation collectors due to adsorption.

As with tropospheric air, the concentration and flux of ⁷Be in rainfall varies seasonally as well (Fig. 2). Maximum fluxes and ecosystem inventories are often recorded in the spring to early summer (Olsen et al. 1985; Brown et al. 1989; Dibb 1989; Bachuber and Bunzl 1992; Baskaran 1995). Brown et al. (1989) and Harvey and Matthews (1989) determined that ⁷Be activities in rainfall can vary by a factor of greater than 20 between individual rainstorms. This suggests that individual storms can potentially dominate ⁷Be deposition. Therefore, to obtain an accurate flux of ⁷Be to the earth's surface, samples must be collected from every storm for at least 1 full year. For investigators who have sampled at least 12 months and acidified their samples in the collectors, annual atmospheric flux of ⁷Be observed from bulk atmospheric deposition collectors has ranged from approximately 1000 to 6500 Bq m⁻² (Turekian et al. 1983; Olsen et al. 1985; Dominik et al. 1987; Schuler et al. 1991; Wallbrink and Murray 1994). Vogler et al. (1996) reported an annual flux at Lake Constance, Germany of 2250 Bq m⁻². Deposition was highest during the summer and related strongly to rainfall (mm d⁻¹). Harvey and Matthews (1989) measured ⁷Be for a year on the Hokitika, South Island, New Zealand. The annual precipitation was about 2.8 m. Wet deposition accounted for about 97% of the total deposition that was 6350 Bq m⁻², and the ⁷Be flux was proportional to the precipitation volume. Dry deposition apparently contributes <10% of ⁷Be inventories (Brown et al. 1989; Harvey and Matthews (1989); Todd and Wong (1989); Walbrink and Murray 1994; Benitez-Nelson and Buesseler 1999; Kaste 1999, Kaste et al. 1999).

⁷Be is delivered to ecosystems primarily as Be⁺² in slightly acidic (pH<6) rainfall. The Be⁺² ion is extremely competitive for cation exchange sites because of its high charge density. As ⁷Be⁺² comes in contact with soils and vegetation, it is rapidly sequestered by exchange surfaces. Forest canopies may decrease the amount of ⁷Be that reaches soils and streams, in contrast to the increases observed for most other analytes (Kaste et al. 1999). After deposition, most ⁷Be decays in the watershed, but some may be exported in particulate or dissolved form.

Table 2. Terrestrial inventories of ^7Be .

Collection date	Location	Approx latitude	Substrate analyzed (n)	Total ^7Be inventory Bq m^{-2}	Source
7/82	Delaware, USA	39°N	marsh + overlying grass	207±13%	Olsen et al. (1985)
7/84	Oak Ridge, TN, USA	36°N	soil + overlying grass	673±3%	Olsen et al. (1985)
1/85	Wallops Island, VA, USA	38°N	marsh + overlying grass	673±7%	Olsen et al. (1985)
1/85	Wallops Island, VA, USA	38°N	unvegetated marsh	107±17%	Olsen et al. (1985)
9/88	Black Mt, Australia	35°S	soil + overlying grass (3)	200±10%	Wallbrink & Murray (1996)
9/88	Black Mt, Australia	35°S	alluvial bare soil (3)	130±20%	Wallbrink & Murray (1996)
5/89	Black Mt, Australia	35°S	soil + overlying grass (3)	400±10%	Wallbrink & Murray (1996)
5/89	Black Mt, Australia	35°S	alluvial bare soil (3)	155±15%	Wallbrink & Murray (1996)
5/96	Idaho, USA	44°N	soil and overlying vegetation (6)	139±16%	Bonniwell et al. (1999)
6/-8/98	Maine, USA	45°N	forested soil (8)	165±40%	Kaste (1999)
12/98	Maine, USA	45°N	bog core + overlying vegetation	554±26%	Kaste (1999)

^7BE DISTRIBUTION IN VEGETATION AND SOILS

^7Be surface inventories (activity of ^7Be per unit area regardless of its vertical distribution) generally range from 100 to 700 Bq m^{-2} . Table 2 summarizes some terrestrial inventories of ^7Be for soil and wetland profiles. Vegetation may be a very important sink for ^7Be , and a significant portion of the ^7Be surface inventory may reside in grasses and forest canopies. Olsen et al. (1985) collected 2 marsh cores in early January 1985 just one meter apart. The vegetated core site had approximately six times the inventory of ^7Be found in the unvegetated core site, with approximately 40% of the ^7Be activity in the grass. Wallbrink and Murray (1996) determined the inventory of ^7Be from soil+grass cores from Black Mountain, Australia from 1988 to 1989. They found ^7Be inventories to double in the same plot in less than one year, and attribute the increase to grass growth (50 to 90 cm increase) which would increase the efficiency of precipitation scavenging. Russell et al. (1981) determined that White Pine (*Pinus strobus* L.) needles from MA, USA contained approximately 1.9 $\text{Bq } ^7\text{Be g}^{-1}$ ash, and found approximately 6 to 9 $\text{Bq } ^7\text{Be g}^{-1}$ ash in twigs from the same tree. They concluded that in a climate with 1.2 meters of rain/year, the forest canopy typically retains 5 to 20% of the open field flux of atmospherically deposited radionuclides. Wallbrink and Murray (1996) calculated that a Eucalypt (*Eucalyptus*) forest canopy retained 4 to 27% of the total ^7Be inventory calculated from fallout measurements. Kaste et al. (1999) reported dried red spruce (*Picea rubens* Sarg.) needles containing approximately 0.15 $\text{Bq } ^7\text{Be g}^{-1}$, and estimated that over 150 $\text{Bq } ^7\text{Be m}^{-2}$ could reside in the forest canopy (ca. 50% of the total surface inventory). ^7Be accumulates over time on vegetation, and is balanced by wash off (precipitation) and decay. Leaves apparently reach steady-state (adsorption = decay) with respect to atmospherically delivered ^7Be before twigs and bark (Fig. 3) (Russell et al. 1981; Norton and Perry, unpublished data 1999).




Figure 3. ^7Be activity on 0 year balsam fir (*Abies balsamea* L. Mill) needles and stems compared with activities on 1 year old balsam fir needles and stems. (Norton and Perry, unpublished data).

^7Be may be a useful tracer for studying erosional processes (Murray et al. 1992). However, such investigations require a detailed understanding of the depth distribution of ^7Be . Wallbrink and Murray (1996) found ^7Be no deeper than 2 cm in bare soil, grassland soil, and eucalyptus forest soil in Australia. Olsen et al. (1985) reported ^7Be was retained in the upper 2 cm of soil in Tennessee, USA and a marsh core collected from Wallops Island, Virginia, USA. However, a second marsh core collected in the summer from Wallops Island had ^7Be as deep as 10 cm. They suggested that the depth of the groundwater table could influence the depth distribution of ^7Be . In central Idaho, Bonniwell et al. (1999) reported that the entire ^7Be inventory was retained in the top 1.2 cm of a soil profile. Kaste (1999) found that the ^7Be residing in the upper 8 cm of the organic horizon at the Bear Brook Watershed in Maine, USA (64 to 300 Bq m⁻²) plus that predicted in the canopy matched the surface inventories predicted by input measurements. Apparently the drainage structure, moisture status of the soil, and overlying vegetation govern the inventory and depth distribution of ^7Be .

^7BE IN FRESHWATERS

You et al. (1989) determined partitioning coefficients ($K_d = [\text{Bq kg}^{-1}]/[\text{Bq L}^{-1}]$) for ^7Be between river water and various substrates in the laboratory, and found that most mud, silt, and clay minerals have a K_d near 10^5 L kg^{-1} under neutral to alkaline conditions ($\text{pH} > 6$; see Fig. 9 in Vesely et al., this volume). They reported that the K_d decreases by four orders of magnitude as pH decreases from 6 to 2, and found no significant effect of DOC on the adsorption of ^7Be . Hawley et al. (1986) and You et al. (1989) both found that K_d varies inversely with the amount of suspended particles in freshwater. Li et al. (1984) suggested that higher concentrations of suspended particles increase particle aggregation, thereby decreasing the effective exchange surface area g⁻¹ of suspended matter. Olsen et al. (1986) and Bonniwell et al. (1999) determined field K_d values for ^7Be in freshwaters to be $>10^4$. Vogler et al. (1996) reported that K_d in the water column of Lake Constance, Germany ranged from $10^{5.3}$ to 10^6 .

^7Be mobility may be enhanced in natural waters by the formation of soluble fluoride (F) and organic acid complexes (Vesely et al. 1989; Vesely et al. this volume). If free F anions are available in the pH range 4.5 to 7, BeF^+ may be a significant fraction of the total Be species in solution. This would decrease the charge density of the total ^7Be in solution, and reduce its adsorption to the solid phase. Below pH 4, aluminum (Al) competition for F decreases Be-F complexes (Vesely et al. 1989). Above pH 7, Be

mobility decreases as hydroxide complexes of Be become significant and Be is adsorbed by sediment and precipitating iron (Fe) and Al hydroxides. Veselý et al. (1989) suggested that Be complexes with fulvic acids may increase Be mobility in surface waters by reducing the fraction of the Be^{+2} in solution

Despite numerous publications documenting the production of ^7Be and the flux to the surface of the earth, there are relatively few studies documenting its mobility and export from watersheds. Dominik et al. (1987) found ^7Be activity in filtered streamwater from Switzerland to be $<1.1 \times 10^{-3} \text{ Bq L}^{-1}$, but found fluvial suspended matter to have an activity of up to 0.49 Bq g^{-1} , averaging 0.132 Bq g^{-1} . Bonniwell et al. (1999) reported that dissolved Be in a mountain stream in central Idaho was below detection limit ($<3 \times 10^{-4} \text{ Bq L}^{-1}$), but suspended sediment generally had 0.05 to 1 Bq g^{-1} . Kaste (1999) found that unfiltered streamwater draining an artificially acidified catchment (pH 4.7) in central Maine had up to 1 Bq L^{-1} during a large snowmelt/rain event. ^7Be was up to 0.75 Bq L^{-1} in unfiltered streamwater from a nearby stream with 20 mg DOC L^{-1} during high discharge. Olsen et al. (1986) reported activity of dissolved ^7Be in the lower Susquehanna and Raritan Rivers eastern USA ranging from approximately 0.002 to 0.003 and 0.002 to 0.014 Bq L^{-1} , respectively. ^7Be activities on suspended particles ranged from ~ 0.26 to 0.44 Bq g^{-1} . In both the Raritan and the Susquehanna, dissolved ^7Be activities (Bq L^{-1}) were higher when the total particle concentration (mg L^{-1}) was higher. In a ^7Be budget for the James River, Virginia, USA, Olsen et al. (1986) determined that less than 5% of the ^7Be in the estuary was from the rivers, and suggested that soils and vegetation in watersheds sequester most of the ^7Be delivered by rainfall. Cooper et al. (1991) studied ^7Be export during a snowmelt event in an arctic watershed. Even though $^{18}\text{O}/^{16}\text{O}$ ratios indicated that $>85\%$ of the discharge during the melt was from the snowpack, over 90% of the ^7Be inventory in the snowpack remained in the watershed after the event.

The efficiency of removal of ^7Be from the water column by particulates may vary among lakes. Krishnaswami et al. (1980) found that the ^7Be inventory in a sediment core from Lake Whitney closely matched the atmospheric flux of ^7Be . They suggested that if sediment focusing effects could be ruled out, the residence time of ^7Be in the lake was very short because of its very high reactivity with particles in the water column. However, in Lake Zurich, Switzerland, Schuler et al. (1991) found that ^7Be fluxes into sediment, measured by sediment traps, accounted for only 30% of the measured atmospheric flux of ^7Be . They proposed that the residence time of ^7Be in the lake was greater than its mean life ($1/\lambda = 76.6$ days). Wan et al. (1987) determined the residence time of ^7Be in Greifensee, Switzerland to be approximately 75 days using calculated atmospheric fluxes and measured ^7Be inventories in sediment. Vogler et al. (1996) observed that the concentration of total ^7Be in Lake Constance, Germany varied with atmospheric input and particle flux. The proportion of ^7Be associated with particles varied with biological productivity, and the residence time calculated with a particle-scavenging model ranged from 40 to 340 days. Steinmann et al. (1999) reported a similar residence time of 50 to 230 days for ^7Be in the water column at Lake Lugano in Switzerland. They found that $\log K_d$ for ^7Be on colloids was approximately 6, while $\log K_d$ on particulates ($>1 \mu\text{m}$) was between 4 and 5. Dominik et al. (1989) calculated the residence time of ^7Be in Lake Geneva to be 60 to 1100 days, using four different methods. They suggested that the most accurate method to determine residence time of ^7Be is to compare its atmospheric flux with ^7Be fluxes in sediment traps. Residence times calculated from ^7Be inventories in bottom sediments were generally within the range of those calculated by sediment traps (400 to 930 days), but the authors warned that small scale heterogeneity could make sediment inventory calculations unreliable.

^7Be inventories from lake sediment cores range from 35 to 875 Bq m^{-2} , with ^7Be

activities in the upper few centimeters of the cores generally ranging from 0.02 to 0.55 Bq g⁻¹ (Krishnaswami et al. 1980; Wan et al. 1987; Dominik et al. 1989). Krishnaswami et al. (1980) found ⁷Be as deep as 3 cm in Lake Whitney, Connecticut, USA, while Wan et al. (1987) reported ⁷Be no deeper than 1 cm in a lake core from Greifensee, Switzerland. Wan et al. (1987) found that ⁷Be inventories in cores ranged by a factor of 4, and were higher where sedimentation rates were higher. Schuler et al. (1991) found a strong seasonality for ⁷Be fluxes in sediment traps in Lake Zurich, with highest fluxes in July and August over three consecutive years. Maximum ⁷Be atmospheric flux and maximum particle flux also occurred during this time period.




Figure 4. Depth-distribution of ⁷Be in the Pacific Ocean. Data Plotted from Silker (1972).

⁷BE IN THE MARINE ENVIRONMENT

Young and Silker (1980) compiled a significant amount of data on dissolved ⁷Be in the Atlantic and Pacific Oceans. They reported that ⁷Be activities in seawater generally ranged from 0.0017 to 0.0117 Bq L⁻¹. Total inventories ranged from approximately 80 to 600 Bq m⁻². They found that concentrations and inventories were highest in areas associated with high rainfall, and increased with latitude. Silker (1972) reported on the depth distribution of ⁷Be in the North Pacific Ocean (Fig. 4), and determined that ⁷Be distribution generally reflected the temperature profile. In cases where there was a strong thermocline, ⁷Be activity was relatively uniform in the mixed (upper) layer, and concentrations decreased rapidly below the thermocline.

Since Young and Silker (1980), several authors have reported ⁷Be activities in seawater (Table 3). Comparing the activities of dissolved ⁷Be and K_d values reported for marine waters with the few data available for freshwater, it appears that ⁷Be can be partitioned favorably into the aqueous phase in the marine environment. Considering the strong pH dependence of Be partition coefficients (You et al. 1989; Vesely et al. 1989; Vesely et al., this volume) the high pH of seawaters relative to many freshwaters should decrease ⁷Be in the aqueous phase. Dibb and Rice (1989a) found that sorption of ⁹Be in laboratory experiments increased with increasing salinity. The wide range of dissolved ⁷Be and K_d values indicates that several variables are likely governing the solubility and partitioning of Be in seawater including pH, salinity, concentrations of suspended matter concentrations (especially Fe) and DOC, competing cations, and the residence time of the suspended load.

Table 3. Dissolved and particulate ^7Be activities in seawater.

Location [USA unless indicated] (time collected)	Dissolved ^7Be (Bq L $^{-1}$)	^7Be on suspended matter [(Bq g $^{-1}$), % of total ^7Be]	log K_d	Suspended load (mg L $^{-1}$)	Source
Sequim Bay, WA	0.0009±50%	0.26±24%, 50	5.4	3.55	Bloom & Crecelius (1983)
Sequim Bay, WA	0.0009±38%	0.28±78%, 24	6.0	1.07	Bloom & Crecelius (1983)
Gironde Estuary, France	0.0004 to .002	0.007-0.41	4.5- 5	n.a.	Martin et al. (1986)
Chesapeake Bay, MD (Spring)	0.013±3% to 0.020±2%	0.41-0.47±2%, 36	4.4- 4.5	11 to 24	Olsen et al. (1986)
Chesapeake Bay, MD (Fall)	0.001±33% to 0.003±14%	0.17±11% to 0.43±4%, 20 to 54	4.8- 5.3	4 to 6	Olsen et al. (1986)
Chesapeake Bay, MD (Year)	<0.0008 to 0.048±2%	<0.007-2.5, 1 to 86	2.8- 6	1 to 121	Dibb & Rice (1989b)
Sabine-Neches Estuary, TX	0.003±15% to 0.035±4%	0.08-0.55±9%, 7 to 57	3.2- 4.9	8 to 92	Baskaran et al. (1997)

Bloom and Crecelius (1983) determined that ^7Be in seawater had a "limited affinity" for suspended matter, and the percent ^7Be adsorbed to particulates is directly proportional to suspended load. Using laboratory experiments with ^7Be and natural "detritus", they demonstrated that high-suspended loads (20 mg L $^{-1}$) had approximately 50% of the total ^7Be adsorbed to suspended matter. At approximately 1 mg L $^{-1}$ suspended matter, only 20% of the total ^7Be in the water column was adsorbed. They also showed that ^7Be adsorbed similarly to detritus and amorphous Fe(OH) $_3$, while adsorption to the algal culture *Pavlova lutheri* (monochrysis) was significantly lower. Although the percent ^7Be adsorbed to suspended matter was directly proportional to suspended load concentrations, the K_d was inversely proportional to suspended load concentrations for their field data (Table 3). Martin et al. (1986) also noted that the K_d was lowest in the area with the maximum suspended load. Li et al. (1984) found that K_d for ^7Be and several other particle-reactive nuclides in seawater was inversely proportional to the suspended load concentrations, in agreement with Hawley et al. (1986) and You et al. (1989) for freshwater systems.

Olsen et al. (1986) found ^7Be activities in filtered river-estuarine and coastal waters along the northeastern USA seaboard ranging from 1.1×10^{-3} to 2×10^{-2} Bq L $^{-1}$, with total (unfiltered) ^7Be reaching 3×10^{-2} Bq L $^{-1}$. Most of the ^7Be was dissolved, typically 60 to 90%. A major factor controlling the concentration of ^7Be on suspended matter was the amount of time the particles remained in the water column. ^7Be activities on suspended particles were highest in high-energy areas, and lowest in low energy areas with high sedimentation rates. Dissolved ^7Be averaged 60% of total ^7Be in Chesapeake Bay (Dibb and Rice 1989a). However, activities of ^7Be on suspended matter reached 2.5 Bq g $^{-1}$, and partition coefficients ranged by over 3 orders of magnitude up to log $K_d = 6$. They concluded, as did Olsen et al. (1986), that rapid variations in the supply of particulates in estuaries and bays may not permit equilibrium to be reached between dissolved ^7Be and the solid phase.

Olsen et al. (1986) determined log K_d for river-estuarine and coastal waters from the northeastern USA coast to range from 3.85 to 5.3, with a median of 4.6. They found a positive correlation between K_d and the Fe concentration of the suspended matter, and suggested that the precipitation of Fe may be an important factor governing the

scavenging of ^7Be . Baskaran et al. (1997) reported relatively low K_d values for ^7Be in the Sabine-Neches estuary, Texas, USA. They found $\log K_d$ to range from 3.18 to 4.94. There was no significant correlation between suspended load and K_d . Baskaran et al. (1997) attributed their low K_d values to the high concentrations of DOC (5 to 20 mg L⁻¹) and suggested that DOC could play a significant role in the fate and the residence times of particle-reactive nuclides such as ^7Be in the water column.

Inventories of ^7Be in near-shore marine sediments range from <30 to 6000 Bq m⁻², with ^7Be activities ranging up to ~0.65 Bq g⁻¹ (Krishnaswami et al. 1980; Olsen et al. 1986; Dibb and Rice 1989b; Canuel et al. 1990, and Summerfield et al. 1999). Maximum inventories of ^7Be in marine sediment commonly occur in the spring, and in areas where fine particles are accumulating rapidly. Canuel et al. (1990) examined the spatial and vertical distribution of ^7Be in sediments of Cape Lookout Bight, NC, USA. They sampled 10 cores in September, 1987, and found ^7Be inventories ranging from approximately 650 to 1375 Bq m⁻² in a <30 m² area. ^7Be penetration reached a maximum depth in the sediment profile during the summer months. ^7Be activity reached 0.033 Bq g⁻¹ 13 cm deep in the core. The authors attributed this seasonal pattern in ^7Be depth distribution to the formation of methane bubble tubes during the summertime. Olsen et al. (1986) found that ^7Be reached a depth of 8 cm in coastal sediments of the eastern USA, but generally was retained in the upper 4 cm. Sneed (1986) determined ^7Be and ^{210}Pb in cores from a dredging site with a high sedimentation rate in New York City harbor. She found ^7Be to a depth of 4 cm in an area where the sedimentation rate was approximately 7 cm yr⁻¹.

APPLICATIONS OF ^7Be

A detailed understanding of the transport and residence time of aerosols in the atmosphere is desired because particles in the micron and sub-micron range probably play a significant role in climate change and trace metal cycling. ^7Be is a very useful tracer in atmospheric transport studies because it (1) has a short half-life, (2) is produced at a relatively constant rate, (3) rapidly attaches to aerosols after formation, and (4) has a large concentration gradient between the stratosphere and troposphere (Bhandari and Rama 1963; Junge 1963; Dutkiewicz and Husain 1979, 1985; Viezee and Singh 1980). In the past decade in particular, ^7Be measurements in the atmosphere have been used to test and validate chemical transport models in the atmosphere (Brost et al. 1991; Rehfeld and Heimann 1995; Koch et al. 1996). Dibb and Jaffrezo (1993) used ^7Be and $^{210}\text{Pb}_{\text{ex}}$ (decay product of ^{222}Rn , $T_{1/2} = 22.3$ yr) to investigate short term "communication" between the atmosphere and the Greenland Ice Sheet. They found that concentrations of ^7Be and ^{210}Pb in fresh snow did not have the spring and fall peaks that ground level aerosol concentrations displayed. On the basis of these data, Dibb and Jaffrezo (1993) suggested that caution must be used when inferring atmospheric chemistry from fresh snow.

The potential for using ^7Be in atmospheric transport studies is particularly great when it is coupled with a nuclide with a contrasting source function and/or a different half-life. The initial production ratio of $^{10}\text{Be}/^7\text{Be}$ is approximately 0.5 (Lal and Peters 1967). As air moves away from the region of maximum production, the $^{10}\text{Be}/^7\text{Be}$ ratio on aerosols in the air mass will increase, because ^7Be has a much shorter radioactive half-life ($T_{1/2} = 53$ days) than ^{10}Be ($T_{1/2} = 1.5$ m.y.). Thus the $^{10}\text{Be}/^7\text{Be}$ ratio in the atmosphere can be used as a "clock" to determine air mass age, and has been used to indicate the intrusion of stratospheric air into the troposphere (Raisback et al. 1981; Dibb et al. 1994; Rehfeld and Heimann 1995; Koch and Rind 1998). Because the residence time of aerosols in the troposphere is fairly short (ca. 1 month [Durana et al. 1996; Bleichrodt 1978]) compared to those in the stratosphere (about 1 year), stratospheric air masses characteristically have a relatively high $^{10}\text{Be}/^7\text{Be}$ ratio. Dibb et al. (1994) found that the $^{10}\text{Be}/^7\text{Be}$ ratio was

nearly constant at 2.2 throughout the year in northwestern Canada. They concluded that a significant portion of the aerosols in the arctic troposphere must be derived from the stratosphere, likely a result of vertical mixing between the two air masses rather than stratospheric injections. Koch and Rind (1998) used a General Circulation Model (GISS GCM) to suggest that the spring maximum in the $^{10}\text{Be}/^{7}\text{Be}$ ratio observed by Dibb et al. (1994) in the lower stratosphere may have resulted from the transport of equatorial air to higher latitude during the previous winter.

The contrasting source terms of ^7Be and ^{210}Pb make the $^7\text{Be}/^{210}\text{Pb}$ ratio a useful tool for investigating air mass sources and vertical transport within the troposphere (Dibb et al. 1992; Baskaran et al. 1993; Rehfeld and Heimann 1995; Koch et al. 1996; Graustein and Turekian 1996, and Benitez-Nelson and Buesseler 1999). While ^7Be is produced primarily in the upper atmosphere, ^{210}Pb is derived from the decay of ^{222}Rn gas ($T_{1/2}=3.8$ days) that is emitted primarily from the surface of the continents. Therefore, ^{210}Pb concentrations in air should decrease with altitude and distance from land. However, ^7Be concentrations in air will be uniform over land and sea, always increasing with altitude. High $^7\text{Be}/^{210}\text{Pb}$ ratios may be an indication of stratosphere-troposphere exchange, or an air mass with components originating from the open ocean. Benitez-Nelson and Buesseler (1999) found that the average $^7\text{Be}/^{210}\text{Pb}$ ratio in rainfall from Woods Hole, MA and Portsmouth, NH (USA) was 12.6 and 20.1, respectively. These data suggest that Woods Hole received air from a more continental source than Portsmouth during the sampling period. The efficiency and rate of vertical transport (convection) within the troposphere will also affect the $^7\text{Be}/^{210}\text{Pb}$ depositional ratio. Increased vertical transport should increase the amount of ^7Be delivered to the ground, and increase the rate at which ^{222}Rn moves upwards. That is, ^7Be is enriched in the lower troposphere while ^{210}Pb is diluted. Baskaran (1995) found that the $^7\text{Be}/^{210}\text{Pb}$ ratio (measured in depositional fluxes) was highest in the summertime (14.4 to 23.9) and lowest in the winter (12.8 to 13.7). He attributed the increased $^7\text{Be}/^{210}\text{Pb}$ ratios during the warmer months to decreased stability of the troposphere (increased convection).

^7Be has useful applications in sedimentological studies. Olsen et al. (1989) used the ratio of short-lived ^7Be to longer-lived excess ^{210}Pb ($^{210}\text{Pb}_{\text{ex}}$) to quantify the amount of re-suspended sedimentary material in coastal waters. ^7Be is very low on bottom sediments relative to its activity on suspended particulates, while $^{210}\text{Pb}_{\text{ex}}$ is nearly the same on bottom sediments and suspended particulates. Using $^7\text{Be}/^{210}\text{Pb}_{\text{ex}}$, they determined that re-suspended bottom sediments may account for more than 80% of the suspended matter in the Savannah Estuary. Canuel et al. (1990) concluded that in coastal sediments unaffected by bioturbation, ^7Be is a useful tracer of sediment accumulation. They found that sediment cores taken before and after storm events can be used to quantify the impacts of individual storms by identifying "new" pulses of sediment. Fitzgerald et al. (2001) used this technique to determine monthly sediment deposition and re-suspension rates in PCB-rich sediments in the Fox River system in Wisconsin, USA. They found that short-term accumulation rates determined by ^7Be were up to 130 times higher than those determined with ^{137}Cs , indicating the dynamic nature of sediment transport in this impounded system. Sommerfield et al. (1999) used ^7Be as a tracer of fine-grained river sediment on the northern California, USA continental margin. They found that ^7Be was present in shelf and slope deposits only after periods of extremely high river discharge, and determined that storms and floods play a significant role in sedimentary processes in modern environments in their study area. ^7Be has been used to calculate sediment accumulation rates in shallow marine environments (Sneed 1986). Canuel et al. (1990) and Dibb and Rice (1989b) found that short-term sediment accumulation rates determined with ^7Be were comparable to longer-term sediment accumulation rates determined from ^{210}Pb . Dibb and Rice (1989b) suggested that the general agreement between the ^7Be and

^{210}Pb derived sedimentation rates could indicate that the processes operating on the Chesapeake Bay sediments have remained relatively constant for the past 100 years. Steinmann et al. (1999) used ^7Be to calculate the coagulation rate of colloids in the epilimnion of Lake Lugano. They measured the sedimentary flux of ^7Be , as well as concentrations of dissolved, colloidal, and particulate ^7Be in the epilimnion. Based on the 2.2 to 16.3 day residence time of ^7Be in the colloidal fraction, they reported coagulation rates of 0.06 to 0.46 d^{-1} .

^{234}Th ($T_{1/2} = 24$ days; derived from decay of ^{238}U) can be used in conjunction with ^7Be to investigate sedimentation processes in areas with a salinity gradient. Both nuclides are particle-reactive, but the source of ^{234}Th is ^{238}U , which is positively correlated with salinity. Feng et al. (1999a, 1999b) utilized the $^{234}\text{Th}_{\text{ex}}/^7\text{Be}$ activity ratios on suspended and bottoms sediments as a tracer of particle sources and transport in the Hudson River estuary, where Th_{ex} is the total Th activity of the sediment minus that supported by ^{238}U . By using the activity ratio of these two nuclides, the apparent effects of grain size, composition, and concentration on adsorption are minimized. They analyzed sediment during different flow regimes and tidal cycles to determine the relative importance of local sediment re-suspension and advection of sediments through the estuary. They found that both radionuclides were removed from the water column on time scales from <1 to 13 days. They also noted that the $^{234}\text{Th}_{\text{ex}}/^7\text{Be}$ tended to be equal or greater than that predicted by local equilibrium (*in situ* production of $^{234}\text{Th}_{\text{ex}}$, atmospheric inputs of ^7Be , scavenging, particles settling and re-suspension) during low flow, which can be explained by transport of particles up-estuary by estuarine circulation. During high flows they observed lower activity ratios, indicating the importance of re-suspension. Feng et al. (1999b) reported that during the course of tidal cycles, the $^{234}\text{Th}_{\text{ex}}/^7\text{Be}$ values were comparable in suspended particles from surface and bottom water with different salinities, which indicated that particles are mixed fairly well in the water column.

The strong affinity of ^7Be for small particles and its steady production make ^7Be a unique tool for hydrologists and geomorphologists. Walling et al. (1999) used ^7Be measurements to document event-based soil redistribution. They found that the spatial patterns of ^7Be and ^{137}Cs (fission product, $T_{1/2} = 30.2$ yr) total inventories were similar. By comparing the inventories of these nuclides to a reference inventory (taken in a stable area where neither erosion or aggradation is occurring) before and after a storm, they could identify areas of soil loss and accumulation on short (a few days) and medium (a few decades) time scales. Bonniwell et al. (1999) used ^7Be and other fallout nuclides to trace sediment and determine erosion rates in a small watershed in Idaho. They used ^7Be as an indicator of "new" sediment during snowmelts, by assuming that "old" sediment that gets re-suspended in the water column would have a negligible concentration of ^7Be . Bonniwell et al. (1999) calculated erosion rates in the watershed by comparing the total export of radionuclides with the measured inventories in the soils.

The different depth distributions of ^7Be , $^{210}\text{Pb}_{\text{ex}}$ and ^{137}Cs (fission product, $T_{1/2} = 30.2$ yr) can be used to fingerprint the soil depth from which suspended sediment was originally removed (Burch et al. 1988; Walling and Woodward 1992; Wallbrink et al. 1999) (Fig. 5). Because ^7Be is often concentrated in the uppermost soil horizons, Murray et al. 1992 suggested using ^7Be as a tracer of topsoil movement. For example, suspended sediment with a high activity of ^7Be (and/or $^{210}\text{Pb}_{\text{ex}}$) could indicate an initial source area at the top of the soil profile, while sediment with a lower activity of ^7Be (and/or $^{210}\text{Pb}_{\text{ex}}$) and high activity of ^{137}Cs could have been derived from deeper in the profile. Wallbrink and Murray (1993) extended the use of these fallout nuclides to identify the type erosion process (sheet flow, rill or gully erosion) removing soil from the landscape (Fig. 6).




Figure 5. Depth-distribution of fallout radionuclides applicable for erosion studies. Generalized from Wallbrink et al. (1999).




Figure 6. Generalized ^7Be and ^{137}Cs labeling of sediment (after Burch et al. 1988). This concept is based on the theory that sediment retains the radionuclide signature originally obtained when it was soil. Used by permission of the editor of *Hydrological Processes*, from Wallbrink and Murray (1993), Fig. 1c, p. 298.

SUMMARY AND CONCLUDING REMARKS

^7Be is a short-lived naturally occurring radionuclide with a unique source term and a relatively constant production rate. After ^7Be is formed in the atmosphere, it rapidly adsorbs to aerosols. Because the residence time of aerosols in the stratosphere is greater than the half-life ^7Be by more than a factor of six, steady-state (production = decay)

equilibrium maintains nearly constant concentrations above the tropopause. Tropospheric concentrations of ^7Be are more dynamic and are generally 1 to 2 orders of magnitude lower than in the stratosphere due to wash out and decay. Near surface air exhibits fluctuations on decadal, semi decadal (4 to 7 years), annual, and semi-annual timescales. The short-term variations of ^7Be concentration in near surface air are a result of variation in production, wash out by precipitation, vertical mixing between and within the stratosphere and troposphere, and horizontal air mass movement/mixing within the troposphere. ^7Be terrestrial inventories and ecosystem fluxes (measured by bulk atmospheric deposition collectors) vary seasonally due to variation in tropospheric ^7Be and precipitation amount. ^7Be is deposited on ecosystems primarily by wet deposition; dry deposition is apparently <10% of the total flux.

^7Be deposited from rainfall rapidly adsorbs to vegetation, soils, and sediment, and in many cases will remain adsorbed to the solid phase until decay. In the soil profile, ^7Be is generally retained in the upper layers. Most of the movement of ^7Be within and from watersheds appears to be in particulate form. The residence time of ^7Be in the water column of lakes is uncertain, reports range from a few weeks to a few months. The partitioning of ^7Be in the marine environment is also poorly understood, with reported K_d values ranging over three orders of magnitude. In estuaries, a significant portion of the ^7Be may be in dissolved form, especially in the presence of DOC. Sediment inventories of ^7Be in both lakes and shallow ocean water exhibit seasonal variations.

^7Be studies have provided scientists with much insight into atmospheric, geochemical, erosional, sedimentological, and nuclear processes. Its decay energy is fairly easy and inexpensive to measure. Therefore some of the more recently developed applications of ^7Be , particularly in atmospheric, erosion, and sedimentation studies, have significant potential for evaluation of environmental problems. However, most of the applications of ^7Be require a very detailed understanding of the distribution and geochemistry of ^7Be , which is still somewhat incomplete in terrestrial ecosystems. In particular, more information is needed on the depth distribution of ^7Be in soils with respect to different drainage conditions, bulk density, and amount of macropore flow channels. Also, the partitioning of ^7Be and its residence time in lakes and the ocean is poorly understood. The effect of competing ligands and DOC on the adsorption behavior needs further documentation, as is evident in the wide range of K_d s reported (three orders of magnitude) in estuaries and tidal bays. With further studies on the geochemical behavior (see Veselý et al., this volume) and distribution of ^7Be under various conditions, it will be possible to take full advantage of this nuclide in earth science investigations.

ACKNOWLEDGMENTS

Our research was supported by the University of Maine and by the National Science Foundation (Grant EAR-9725705). In addition, this review paper was greatly improved by the comments of Peter Wallbrink, Curtis Olsen, and Jack Dibb.

REFERENCES

- Ajzenberg-Selove F (1988) Energy levels of the light nuclei $A = 5-10$. Nuclear Physics A 490:1-225
- Arnold JR (1956) Beryllium-10 produced by cosmic rays. Science 124:584-585
- Arnold JR, Al-Salih H (1955) Beryllium-7 produced by cosmic rays. Science 121:451-453
- Bachhuber H, Bunzl K (1992) Background levels of atmospheric deposition to ground and temporal variation of ^{129}I , ^{127}I , ^{137}Cs , and ^7Be in a rural area of Germany. J Environ Radioact 16:77-89
- Baskaran M (1995) A search for the seasonal variability on the depositional fluxes of ^7Be and ^{210}Pb . J Geophys Res 100:2833-2840
- Baskaran M, Coleman CH, Santschi PH (1993) Atmospheric depositional fluxes of ^7Be and ^{210}Pb at Galveston and College Station, Texas. J Geophys Res 98:20555-20571

- Baskaran M, Ravichandran M, Bianchi TS (1997) Cycling of ^7Be and ^{210}Pb in a high DOC, shallow, turbid estuary of southeast Texas. *Estuarine Coastal Shelf Sci* 45:165-176
- Beer J, Blinov A, Bonani G, Finkel RC, Hofmann HJ, Lehmann B, Oeschger H, Sigg A, Schwander J, Staffelbach T, Stauffer B, Suter M, Wolfli W (1990) Use of ^{10}Be in polar ice to trace the 11-year cycle of solar activity. *Nature* 347:164-166
- Benitez-Nelson C, Buesseler KO (1999) Phosphorous-32, phosphorous-37, beryllium-7, and lead-210: Atmospheric fluxes and utility in tracing stratosphere/troposphere exchange. *J Geophys Res* 104:11745-11754
- Bhandari N, Rama (1963) Atmospheric circulation from observations of sodium-22 and other short-lived natural radioactivities. *J Geophys Res* 68:1959-1966
- Bhandari N, Lal, D, Rama (1970) Vertical structure of the troposphere as revealed by radioactive tracer studies. *J Geophys Res* 75:2974-2980
- Bleichrodt JF (1978) Mean tropospheric residence time of cosmic-ray-produced beryllium-7 at north temperate latitudes. *J Geophys Res* 83:3058-3062
- Bloom N, Crecelius EA (1983) Solubility behavior of atmospheric ^7Be in the marine environment. *Marine Chem* 12:323-331
- Bonniwell EC, Matisoff G, Whiting PJ (1999) Determining the times and distances of particle transit in a mountain stream using fallout radionuclides. *Geomorph* 27:75-92
- Bouchez PR, Tobailem J, Robert J, Muxart R, Mellet R, Daudel P, Daudel R (1956) Nouvelle détermination de la différence des périodes de ^7Be métallique et de BeF_2 . *J Phys Radium* 17:363-365
- Brost RA, Feichter J, Heimann M (1991) Three-dimensional simulation of ^7Be in a global climate model. *J Geophys Res* 96:22423-22445
- Brown L, Stensland GJ, Klein J, Middleton R (1989) Atmospheric deposition of ^7Be and ^{10}Be . *Geochim Cosmochim Acta* 53:135-142
- Burch GJ, Barnes CJ, Moore ID, Barling RD, Mackenzie DJ, Olley JM (1988) Detection and prediction of sediment sources in catchments: Use of ^7Be and ^{137}Cs . *In Hydrology and Water Resources Symposium. Aust Inst Eng National Conf Publ* 88:146-151
- Canuel EA, Martens CS, Benninger LK (1990) Seasonal variations in the ^7Be activity in the sediments of Cape Lookout Bight, North Carolina. *Geochim Cosmochim Acta* 54:237-245
- Cooper LW, Olsen CR, Solomon DK, Larsen IL, Cook RB, Grebmeier JM (1991) Stable isotopes of oxygen and natural and fallout radionuclides used for tracing runoff during snowmelt in an arctic watershed. *Water Resources Res* 27:2171-2179
- Dibb JE (1989) Atmospheric deposition of beryllium-7 in the Chesapeake Bay Region. *J Geophys Res* 94:2261-2265
- Dibb JE, Jaffrezo, JL (1993) Beryllium-7 and lead-210 in aerosol and snow in the Dye 3 Gas, Aerosol, and Snow Sampling Program. *Atmos Environ* 27A:2751-2760
- Dibb JE, Rice DL (1989a) The geochemistry of beryllium-7 in Chesapeake Bay. *Estuarine, Coastal, Shelf Sci* 28:379-394
- Dibb JE, Rice DL (1989b) Temporal and spatial distribution of beryllium-7 in the sediments of Chesapeake Bay. *Estuarine Coastal Shelf Sci* 28:395-406
- Dibb JE, Talbot RW, Gregory GL (1992) Beryllium-7 and lead-210 in the western arctic atmosphere: Observations from three recent aircraft-based sampling programs. *J Geophys Res* 97:16709-16715
- Dibb JE, Meeker DL, Finkel RC, Southon JR, Caffee MW (1994) Estimation of stratospheric input to the arctic troposphere: ^7Be and ^{10}Be in aerosols at Alberta, Canada. *J Geophys Res* 99:12855-12864
- Dingle AN (1965) Stratospheric tapping by intense convective storms: Implications for public health in the United States. *Science* 148:227-229
- Dominik J, Burrus D, Vernet J-P (1987) Transport of the environmental radionuclides in an alpine watershed. *Earth Planet Sci Lett* 84:165-180
- Dominik J, Schuler C, Santschi PH (1989) Residence times of ^{234}Th and ^7Be in Lake Geneva. *Earth Planet Sci Lett* 93:345-358
- Dueñas C, Fernández MC, Liger E, Carretero J (1999) Gross alpha, gross beta activities and ^7Be concentrations in surface air: Analysis of their variations and prediction model. *Atmos Environ* 33:3705-3715
- Durana L, Chudy M, Masarik J (1996) Investigation of ^7Be in the Bratislava atmosphere. *J Radioanal Nucl Chem* 207:345-356
- Dutkiewicz VA, Husain L (1979) Determination of stratospheric and tropospheric components of ^7Be in surface air. *Geophys Res Lett* 6:171-174
- Dutkiewicz VA, Husain L (1985) Stratospheric and tropospheric components of ^7Be in surface air. *J Geophys Res* 90:5783-5788
- Feely HW, Larsen RJ, Sanderson CG (1989) Factors that cause seasonal variations in beryllium-7 concentrations in surface air. *J Environ Radioact* 9:223-249

- Feng H, Cochran JK, Hirschberg DJ (1999a) ^{234}Th and ^7Be as tracers for the transport and dynamics of suspended particles in a partially mixed estuary. *Geochem Cosmochim Acta* 63:2487-2505
- Feng H, Cochran JK, Hirschberg DJ (1999b) ^{234}Th and ^7Be as tracers for the sources of particles to the turbidity maximum of the hudson river estuary. *Estuarine Coastal Shelf Sci* 49:629-645
- Fitzgerald SA, Val Klump J, Swarzenski PW, Mackenzie RA, Richards KD (2001) Beryllium-7 as a tracer of short-term sediment deposition and resuspension in the Fox River, Wisconsin. *Environ Sci Technol* 35:300-305
- Goel PS, Jha S, Lal D, Radhakrishna PR (1956) Cosmic ray produced beryllium isotopes in rain water. *Nuclear Phys* 1:196-201
- Graustein WC, Turekian KK (1996) ^7Be and ^{210}Pb indicate an upper troposphere source for elevated ozone in the summertime subtropical free troposphere of the eastern North Atlantic. *Geophys Res Lett* 23:539-542
- Harvey MJ, Matthews KM (1989) ^7Be deposition in a high-rainfall area of New Zealand. *J Atmos Chem* 8:299-306
- Hawley N, Robbins JA, Eadie BJ (1986) The partitioning of ^7Be in fresh water. *Geochim Cosmochim Acta* 50:1127-1131
- Hensley WK, Bassett WA, Huizenga JR (1973) Pressure dependence of the radioactive decay constant of beryllium-7. *Science* 181:1164-1165
- Hötzl H, Rosner G, Winkler R (1991) Correlation of ^7Be concentrations in surface air and precipitation with the solar cycle. *Naturwissenschaften* 78:215-217
- Huh C-A (1999) Dependence of the decay rate of ^7Be on chemical forms. *Earth Planet Sci Lett* 171:325-328
- Husain L, Coffey PE, Meyers RE, Cederwall RT (1977) Ozone transport from stratosphere to troposphere. *Geophys Res Lett* 4:363-365
- Jaeger M, Wilmes S, Kolle V, Staudt G, Mohr P (1996) Precision measurements of the half-life of ^7Be . *Phys Rev* 54:423-424
- Johlige HW, Aumann DC, Born H-J (1970) Determination of the relative electron density at the Be nucleus in different chemical combinations, measured as changes in the electron-capture half-life of ^7Be . *Phys Rev* 2:1616-1621
- Junge CE (1963) Studies of global exchange processes in the atmosphere by natural and artificial tracers. *J Geophys Res* 68:3849-3856
- Kaste JM (1999) Dynamics of cosmogenic and bedrock-derived beryllium nuclides in forested ecosystems in Maine, U.S.A. Unpublished MSc thesis, University of Maine
- Kaste JM, Norton SA, Fernandez IJ, Hess CT (1999) Delivery of cosmogenic beryllium-7 to forested ecosystems in Maine, USA. *Geol Soc Am Abstr* 31:A305
- Koch DM, Mann, ME (1996) Spatial and temporal variability of ^7Be surface concentrations. *Tellus* 48B:387-396
- Koch DM, Rind D (1998) Beryllium-10/beryllium-7 as a tracer of stratospheric transport. *J Geophys Res* 103:3907-3917
- Koch DM, Jacob DJ, Graustein WC (1996) Vertical transport of aerosols as indicated by ^7Be and ^{210}Pb in a chemical tracer model. *J Geophys Res* 101:18651-18666
- Kostadinov KN, Yanev YL, Mavrodiev VM (1988) A method for the preconcentration of cosmogenic beryllium-7 in natural waters. *J Radioan Nuclear Chem Articles* 121:509-513
- Kraushaar JJ, Wilson ED, Bainbridge KT (1953) Comparison of the values of the disintegration constant of ^7Be in Be, BeO, and BeF₂. *Phys Rev* 90:610-614
- Krishnaswami S, Benninger LK, Aller RC, Damm KL (1980) Atmospherically-derived radionuclides as tracers of sediment mixing and accumulation in near-shore marine and lake sediments: Evidence from ^7Be , ^{210}Pb , and $^{239,240}\text{Pu}$. *Earth Planet Sci Lett* 47:307-318
- Lal D, Peters B (1967) Cosmic ray produced radioactivity on earth. In *Handbuch der Physik* 46:551-612, Sitte K (ed) Springer-Verlag, New York
- Lal D, Malhotra PK, Peters B (1958) On the production of radioisotopes in the atmosphere by cosmic radiation and their application to meteorology. *J Atmos Terrest Phys* 12:306-328
- Larsen IL, Cutshall NH (1981) Direct determination of ^7Be in sediments. *Earth Planet Sci Lett* 54:379-384
- Leininger RF, Segrè E, Wiegand C (1949) Experiments on the effect of atomic electrons on the decay constant of ^7Be . *Phys Rev* 76:897-898
- Leininger RF, Segrè E, Wiegand C (1951) Erratum: Experiments on the effect of atomic electrons on the decay constant of ^7Be . *Phys Rev* 81:280
- Li Y-H, Burkhardt L, Buchholtz M, O'Hara P, Santschi P (1984) Partition of radiotracers between suspended particles and seawater. *Geochim Cosmochim Acta* 48:2011-2019
- Liu LG, Huh CA (2000) Effect of pressure on the decay rate of ^7Be . *Earth Planetary Sci Lett* 180:163-167.

- Martin JM, Mouchel JM, Thomas AJ (1986) Time concepts in hydrodynamic systems with an application to ^7Be in the Gironde Estuary. *Marine Chem* 18:369-392
- Murray AS, Olley JM, Wallbrink PJ (1992) Natural radionuclide behavior in the fluvial environment. *Radiat Prot Dosim* 45:285-288
- Noyce JR, Chen, TS, Moore DT, Beck JN, Kuroda PK (1971) Temporal distributions of radioactivity and $^{87}\text{Sr}/^{86}\text{Sr}$ ratios during rainstorms. *J Geophys Res* 76:646-656
- Olsen C, Larsen IL, Lowry PD, Cutshall NH, Todd JF, Wong GTF, Casey WH (1985) Atmospheric fluxes and marsh-soil inventories of ^7Be and ^{210}Pb . *J Geophys Res* 90:10487-10495
- Olsen CR, Larsen IL, Lowry PD, Cutshall NH (1986) Geochemistry and deposition of ^7Be in river-estuarine and coastal waters. *J Geophys Res* 91:896-908
- Olsen CR, Thein M, Larsen IL, Lowry PD, Mulholland PJ, Cutshall NH, Byrd JT, Windom, HL (1989) Plutonium, lead-210, and carbon isotopes in the Savannah Estuary: Riverborne versus marine sources. *Environ Sci Technol* 23:1475-1481
- Papastefanou C (1991) High-level ^7Be concentrations in air after the SN 1987: A supernova outburst in the large magellanic cloud on 23 February 1987. *Nucl Geophys* 5:51-52
- Peters B (1955) Radioactive beryllium in the atmosphere and on the earth. *Proc Indian Acad Sci* 41:67-71
- Raisbeck GM, Yiou F, Fruneau M, Loiseux JM, Lieuvin M, Ravel JC (1981) Cosmogenic $^{10}\text{Be}/^7\text{Be}$ as a probe of atmospheric transport processes. *Geophys Res Lett* 8:1015-1018
- Rama, Honda M (1961) Natural radioactivity in the atmosphere. *J Geophys Res* 66:3227-3231
- Rehfeld S, Heimann H (1995) Three dimensional atmospheric transport simulation of the radioactive tracers ^{210}Pb , ^7Be , ^{10}Be , and ^{90}Sr . *J Geophys Res* 100:26141-26161
- Reiter ER (1975), Stratospheric-tropospheric exchange processes. *Rev Geophys Space Phys* 13:459-474
- Russell IJ, Choquette CE, Fang S-L, Dundulis WP, Pao AA, Pszeny AAP (1981) Forest vegetation as a sink for atmospheric particulates: Quantitative studies in rain and dry deposition. *J Geophys Res* 86:5247-5363
- Schuler C, Wieland E, Santschi PH, Sturm M, Lueck A, Bollhalder S, Beer J, Bonani G, Hofmann HJ, Suter M, Wolfli W (1991) A multitracer study of radionuclides in Lake Zürich, Switzerland I. Comparison of atmospheric and sedimentary fluxes of ^7Be , ^{10}Be , ^{210}Pb , ^{210}Po , and ^{137}Cs . *J Geophys Res* 96:17051-17065
- Schumann G, Stoepller M (1963) Beryllium 7 in the atmosphere. *J Geophys Res* 68:3827-3830
- Segrè E (1947) Possibility of altering the decay rate of a radioactive substance. *Phys Rev* 71:274-275
- Segrè E, Wiegand CE (1951) Erratum: Experiments on the effect of atomic electrons on the decay constant of ^7Be . *Phys Rev* 81:284
- Silker WB (1972) Horizontal and vertical distributions of radionuclides in the North Pacific Ocean. *J Geophys Res* 77:1061-1070
- Sneed SB (1986) Sediment chronologies in New York Harbor Borrow Pits. Unpublished MSc thesis, State University of New York, Stony Brook
- Sommerfield CK, Nittrouer CA, Alexander CR (1999) ^7Be a tracer of flood sedimentation on the northern California continental margin. *Cont Shelf Res* 19:355-361
- Steinmann P, Billen T, Loizeau JL, Dominik J (1999) Beryllium-7 as a tracer to study mechanisms and rates of metal scavenging from lake surface waters. *Geochem Cosmochim Acta* 63:1621-1633
- Todd JF, Wong TF (1989) Atmospheric depositional characteristics of beryllium 7 and lead 210 along the southeastern Virginia coast. *J Geophys Res* 94:11106-11116
- Turekian KK, Benninger LK, Dion EP (1983) ^7Be and ^{210}Pb total deposition fluxes at New Haven, Connecticut, and at Bermuda. *J Geophys Res* 88:5411-5415
- Vesely J, Benes P, Sevcik K (1989) Occurrence and speciation of beryllium in acidified freshwaters. *Water Resources Res* 23:711-717
- Vieze W, Singh HB (1980) The distribution of ^7Be in the troposphere: Implications on stratospheric/tropospheric air exchange. *Geophys Res Lett* 7:805-808
- Vogler S, Jung M, Mangini (1996) Scavenging of ^{234}Th and ^7Be in Lake Constance. *Limnol Oceanogr* 41:1384-1393
- Wallbrink PJ, Murray AS (1993) Use of fallout radionuclides as indicators of erosion processes. *Hydro Process* 7:297-304
- Wallbrink PJ, Murray AS (1994) Fallout of ^7Be in south Eastern Australia. *J Environ Radioact* 25:213-228
- Wallbrink PJ, Murray AS (1996) Distribution and variability of ^7Be in soils under different surface cover conditions and its potential for describing soil redistribution processes. *Water Resources Res* 32:467-476
- Wallbrink PJ, Murray AS, Olley JM (1999) Relating suspended sediment to its original soil depth using fallout radionuclides. *Soil Sci Soc Am J* 63:369-378

- Walling DE, Woodward JC (1992) Use of radiometric fingerprints to derive information on suspended sediment sources. *In* Erosion and Sediment Transport Monitoring Programmes in River Basins. Int'l Assoc Hydrological Sci Publ 210:153-164
- Walling DE, He Q, Blake W (1999) Use of ^7Be and ^{137}Cs measurements to document short- and medium-term rates of water-induced erosion on agricultural land. *Water Resources Res* 35:3865-3874
- Walton A, Fried RE (1962) The deposition of beryllium-7 and phosphorous-32 in precipitation at north temperate latitudes. *J Geophys Res* 67:5335-5340
- Wan G, Santschi PH, Sturm M, Farrenkothen K, Lueck A, Werth E, Schuler C (1987) Natural (^{210}Pb , ^7Be) and fallout (^{137}Cs , $^{239,240}\text{Pu}$, ^{90}Sr) radionuclides as geochemical tracers of sedimentation in Greifensee, Switzerland. *Chem. Geology* 63:181-196
- You CF, Lee T, Li YH (1989) The partition of Be between soil and water. *Chem. Geology* 77:105-118
- Young J, Silker W (1980) Aerosol depositional velocities on the Pacific and Atlantic Oceans calculated from ^7Be measurements. *Earth Planet Sci Lett* 50:92-104
- Zanis P, Schuepbach E, Gaggeler HW, Hübener S, Tobler L (1999) Factors controlling beryllium-7 at Jungfrauoch in Switzerland. *Tellus* 51B:789-805